



# Quasilinear Elliptic Hamilton-Jacobi Equations on Complete Manifolds

Marie-Françoise Bidaut-Veron, Marta Garcia-Huidobro, Laurent Veron

## ► To cite this version:

Marie-Françoise Bidaut-Veron, Marta Garcia-Huidobro, Laurent Veron. Quasilinear Elliptic Hamilton-Jacobi Equations on Complete Manifolds. 2013. hal-00827016v3

**HAL Id: hal-00827016**

**<https://hal.science/hal-00827016v3>**

Preprint submitted on 4 Jun 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## QUASILINEAR ELLIPTIC HAMILTON-JACOBI EQUATIONS ON COMPLETE MANIFOLDS

Marie-Françoise Bidaut-Véron<sup>1</sup>, Marta Garcia-Huidobro<sup>2</sup>, Laurent Véron<sup>3</sup>

**RÉSUMÉ.** Let  $(M^n, g)$  be a  $n$ -dimensional complete, non-compact and connected Riemannian manifold, with Ricci tensor  $Ricc_g$  and sectional curvature  $Sec_g$ . Assume  $Ricc_g \geq (1-n)B^2$ , and either  $p > 2$  and  $Sec_g(x) = o(dist^2(x, a))$  when  $dist^2(x, a) \rightarrow \infty$  for  $a \in M$ , or  $1 < p < 2$  and  $Sec_g(x) \leq 0$ . If  $q > p - 1 > 0$ , any  $C^1$  solution of (E)  $-\Delta_p u + |\nabla u|^q = 0$  on  $M$  satisfies  $|\nabla u(x)| \leq c_{n,p,q} B^{\frac{1}{q+1-p}}$  for some constant  $c_{n,p,q} > 0$ . As a consequence there exists  $c_{n,p} > 0$  such that any positive  $p$ -harmonic function  $v$  on  $M$  satisfies  $v(a)e^{-c_{n,p} B dist(x,a)} \leq v(x) \leq v(a)e^{c_{n,p} B dist(x,a)}$  for any  $(a, x) \in M \times M$ .

### Equations de Hamilton-Jacobi quasilinéaires sur une variété complète

**RÉSUMÉ.** Soit  $(M^n, g)$  une variété riemannienne  $n$ -dimensionnelle complète, non compacte et connexe de courbures de Ricci  $Ricc_g$  et sectionnelle  $Sec_g$ . On suppose  $Ricc_g \geq (1-n)B^2$  et  $Sec_g(x) = o(dist^2(x, a))$  si  $dist^2(x, a) \rightarrow \infty$  pour  $a \in M$  si  $p > 2$ , ou  $Sec_g(x) \leq 0$  si  $1 < p < 2$ . Si  $q > p - 1 > 0$ , toute solution de classe  $C^1$  de (E)  $-\Delta_p u + |\nabla u|^q = 0$  sur  $M$  satisfait à  $|\nabla u(x)| \leq c_{n,p,q} B^{\frac{1}{q+1-p}}$  où  $c_{n,p,q} > 0$  est une constante. On en déduit qu'il existe  $c_{n,p} > 0$  tel que toute fonction  $p$ -harmonique positive  $v$  sur  $M$  satisfait à l'encadrement suivant,  $v(a)e^{-c_{n,p} B dist(x,a)} \leq v(x) \leq v(a)e^{c_{n,p} B dist(x,a)}$  pour tout  $(a, x) \in M \times M$ .

**Version française abrégée.** Soit  $(M^n, g)$  une variété riemannienne complète, non-compacte et connexe de courbure de Ricci  $Ricc_g$  et courbure sectionnelle  $Sec_g$ . Pour tout  $p > 1$ , on dénote par  $u \mapsto \Delta_p u := \operatorname{div}(|\nabla u|^{p-2} \nabla u)$  le  $p$ -Laplacien sur  $M$  pour la métrique  $g$ . Notre résultat principal est le suivant

**Théorème 1.** Soit  $B \geq 0$  tel que  $Ricc_g \geq (1-n)B^2$  et  $q > p - 1 > 0$ . On suppose

$$(1) \quad \lim_{dist(x,a) \rightarrow \infty} \frac{Sec_g(x)}{(dist(x,a))^2} = 0$$

pour tout  $a \in M$  si  $p > 2$ , ou  $Sec_g \leq 0$  si  $1 < p < 2$ . Il existe alors  $c_{n,p,q} > 0$  telle que toute solution  $u \in C^1(M)$  de

$$(2) \quad -\Delta_p u + |\nabla u|^q = 0 \quad \text{sur } M$$

vérifie

$$(3) \quad |\nabla u(x)| \leq c_{n,p,q} B^{\frac{1}{q+1-p}} \quad \forall x \in M.$$

Une des conséquences est un théorème de type Liouville.

1. Laboratoire de Mathématiques et Physique Théorique, CNRS UMR 7350, Faculté des Sciences, 37200 Tours France. E-mail : veronmf@univ-tours.fr

2. Departamento de Matematicas, Pontifica Universidad Catolica de Chile Casilla 307, Correo 2, Santiago de Chile. E-mail : mgarcia@mat.puc.cl

3. Laboratoire de Mathématiques et Physique Théorique, CNRS UMR 7350, Faculté des Sciences, 37200 Tours France. E-mail : veronl@univ-tours.fr

**Corollaire 2.** *Supposons que  $Ricc_g \geq 0$ ,  $q > p - 1 > 0$  et que les hypothèses du Théorème 1 portant sur la courbure sectionnelle soient vérifiées si  $p \neq 2$ . Alors toute solution  $u \in C^1(M)$  de (2) est constante.*

Si  $v$  est une fonction  $p$ -harmonique positive sur  $M$ , la fonction  $u := -(p - 1) \ln v$  vérifie

$$(4) \quad -\Delta_p u + |\nabla u|^p = 0 \quad \text{sur } M.$$

En utilisant le résultat du théorème 1, on en déduit

**Théorème 3.** *Supposons que  $p > 1$  et que les hypothèses du Théorème 1 portant sur la courbure soient vérifiées. Il existe alors une constante  $c_{n,p} > 0$  telle que toute fonction  $p$ -harmonique et positive  $v$  sur  $M$  vérifie*

$$(5) \quad v(a)e^{-c(n,p)B \text{dist}(x,a)} \leq v(x) \leq v(a)e^{c(n,p)B \text{dist}(x,a)} \quad \forall (a, x) \in M \times M.$$

Quand  $p = 2$  Cheng et Yau [1] ont montré que toute fonction harmonique positive sur une variété riemannienne complète à courbure de Ricci positive est une constante. Dans le cas des fonctions  $p$ -harmoniques positives et sous l'hypothèse de minoration uniforme de la courbure sectionnelle,  $Sec_g \geq -B^2$ , Kotschwar et Ni [4] montrent que toute fonction  $p$ -harmonique positive  $v$  sur  $M$  vérifie l'estimation suivante,

$$(6) \quad \frac{|\nabla v|}{v} \leq (p - 1)B.$$

Notons que leur hypothèse implique  $Ricc_g \geq (1 - n)B^2$ .

---

Let  $(M^n, g)$  be a complete, connected and non compact Riemannian manifold with Ricci curvature  $Ricc_g$  and sectional curvature  $Sec_g$ . For  $p > 1$  we denote by  $\Delta_p$  the  $p$ -Laplacian defined in the metric  $g$  by

$$\Delta_p u := \text{div} \left( |\nabla u|^{p-2} \nabla u \right),$$

and thus  $\Delta_2$  is the Laplace-Beltrami operator on  $M$ . If  $p = 2$  a classical result due to Cheng and Yau [1] asserts that if  $Ricc_g$  is nonnegative, any nonnegative harmonic function  $v$  is a constant. In [4], Kotschwar et Ni obtained sharper results dealing with positive  $p$ -harmonic functions under the assumption that  $Sec_g \geq -B^2$ . They proved that if  $v$  is such a function, it satisfies

$$(1) \quad \frac{|\nabla v|}{v} \leq (p - 1)B.$$

Their assumption on  $Sec_g$  implies  $Ricc_g \geq (1 - n)B^2$ . They also noticed that if  $p = 2$  their estimate holds under the previous lower estimate on the Ricci curvature. In this note we give an extension of their result in imbedding it the more general class of quasilinear Hamilton-Jacobi type equations

$$(2) \quad -\Delta_p u + |\nabla u|^q = 0 \quad \text{on } M.$$

Our main result is the following

**Theorem 1.** *Let  $B \geq 0$  such that  $\text{Ric}_g \geq (1-n)B^2$ . If  $p > 2$  we assume that for any  $a \in M$*

$$(3) \quad \lim_{\text{dist}(x,a) \rightarrow \infty} \frac{\text{Sec}_g(x)}{(\text{dist}(x,a))^2} = 0,$$

*and if  $1 < p < 2$  that  $\text{Sec}_g \leq 0$ . Then there exists  $c_{n,p,q} > 0$  such that any solution  $u \in C^1(M)$  of (2) satisfies*

$$(4) \quad |\nabla u(x)| \leq c_{n,p,q} B^{\frac{1}{q+1-p}} \quad \forall x \in M.$$

A clear consequence of (3) is the following Liouville theorem

**Corollary 2.** *Assume  $\text{Ric}_g \geq 0$  and that the assumptions of Theorem 1 concerning  $\text{Sec}_g$  hold if  $p \neq 2$ . Then any solution  $u \in C^1(M)$  of (2) is constant.*

If  $v$  is a positive  $p$ -harmonic function on  $M$ , then  $u := -(p-1) \ln v$  satisfies

$$(5) \quad -\Delta_p u + |\nabla u|^p = 0 \quad \text{on } M.$$

Therefore estimate (4) yields to the following result

**Theorem 3.** *Assume that  $p > 1$  and the curvature assumptions of Theorem 1 are fulfilled. Then there exists a constant  $c_{n,p} > 0$  such that any positive  $p$ -harmonic function  $v$  on  $M$  satisfies*

$$(6) \quad v(a)e^{-c(n,p)B \text{dist}(x,a)} \leq v(x) \leq v(a)e^{c(n,p)B \text{dist}(x,a)} \quad \forall (a,x) \in M \times M.$$

**Proof of Theorem 1.** Let  $M_+ := \{x \in M : |\nabla u(x)| > 0\}$ . Then  $M_+$  is open and  $u \in C^3(M_+)$  since the equation is no longer degenerate. The proof is based upon the fact that  $z = |\nabla u|^2$  is a subsolution of an elliptic differential inequality with a superlinear absorption term (see [5] for other applications). We denote by  $TM$  the tangent bundle of  $M$  and by  $\langle \cdot, \cdot \rangle$  the scalar product induced by the metric  $g$ . We recall that any  $C^3$ -function  $u$  verifies the Böchner-Weitzenböck formula; combined with Schwarz inequality it yields to

$$(7) \quad \begin{aligned} \frac{1}{2} \Delta_2 |\nabla u|^2 &= |D^2 u|^2 + \langle \nabla \Delta_2 u, \nabla u \rangle + \text{Ric}_g(\nabla u, \nabla u) \\ &\geq \frac{1}{n} |\Delta_2 u|^2 + \langle \nabla \Delta_2 u, \nabla u \rangle + \text{Ric}_g(\nabla u, \nabla u), \end{aligned}$$

where  $D^2 u$  is the Hessian. If  $u$  is a  $C^1$  solution of (2), then  $z = |\nabla u|^2$  satisfies

$$(8) \quad -\Delta_2 u - \frac{p-2}{2} \frac{\langle \nabla z, \nabla u \rangle}{z} + z^{\frac{q+2-p}{2}} = 0$$

on  $M_+$ . Replacing  $\Delta_2 u$  in (7) it follows that, for any  $a > 0$ ,

$$(9) \quad \begin{aligned} \Delta_2 z + (p-2) \frac{\langle D^2 z(\nabla u), \nabla u \rangle}{z} &\geq \frac{2a^2}{N} z^{q+2-p} - \frac{1}{Na^2} \frac{\langle \nabla z, \nabla u \rangle^2}{z^2} - \frac{(p-2)}{2} \frac{|\nabla z|^2}{z} \\ &\quad + (p-2) \frac{\langle \nabla z, \nabla u \rangle^2}{z^2} + (q+2-p) z^{\frac{q-p}{2}} \langle \nabla z, \nabla u \rangle - (N-1)B^2 z. \end{aligned}$$

Since  $z^{\frac{q-p}{2}} |\langle \nabla z, \nabla u \rangle| \leq z^{\frac{q+1-p}{2}} \frac{|\nabla z|}{\sqrt{z}}$ , we can take  $a = a(p,q) > 0$  large enough so

that the right-hand side of (9) is bounded from below by  $Cz^{q+2-p} - D \frac{|\nabla z|^2}{z}$  for

some  $C, D > 0$  which depend only on  $p$  and  $q$ . We set

$$\mathcal{A}(v) := -\Delta_2 v - (p-2) \frac{\langle D^2 v(\nabla u), \nabla u \rangle}{|\nabla u|^2} = - \sum_{i,j=1}^N a_{ij} v_{x_i x_j}$$

where the  $a_{ij}$  depend on  $\nabla u$  and satisfy

$$\theta |\xi|^2 \leq \sum_{i,j=1}^N a_{ij} \xi_i \xi_j \leq \Theta |\xi|^2 \quad \forall \xi = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n,$$

where  $\theta = \min\{1, p-1\}$  and  $\Theta = \max\{1, p-1\}$ . Then

$$(10) \quad \mathcal{L}^*(z) := \mathcal{A}(z) + C z^{q+2-p} - D \frac{|\nabla z|^2}{z} - (n-1)B^2 z \leq 0 \quad \text{in } M_+.$$

The next lemma is a local estimate.

**Lemma 1.** *Let  $B_R(a) \subset M^n(g)$  be the geodesic ball of radius  $R > 0$  and center  $a$ . Assume that  $\text{Ric}_g \geq -(n-1)B^2$  and either  $\text{Sec}_g \geq -S^2$  for some  $S^2 := S_R^2$  in  $B_R(a)$  if  $p > 2$ , or  $\text{Sec}_g \leq 0$  if  $1 < p < 2$ . Then there exists  $c = c(n, p, q) > 0$  such that the function*

$$(11) \quad w(x) = \lambda (R^2 - r^2(x))^{-\frac{2}{q+1-p}} + \mu \quad \text{with } r = r(x) = d(x, a),$$

satisfies  $\mathcal{L}^*(w) \geq 0$  in  $B_R(a)$ , provided

$$(12) \quad \lambda = c \max \left\{ (R^4 B^2)^{\frac{1}{q+1-p}}, ((1+B+(p-2)_+ S) R^3)^{\frac{1}{q+1-p}} \right\}$$

and

$$(13) \quad \mu = ((n-1)B^2)^{\frac{1}{q+1-p}}.$$

**Proof.** We recall that  $\Delta_2 w = w'' + w' \Delta_2 r$  and by [6, Lemma 1]

$$\Delta_2 r \leq (n-1)B \coth(Br) \leq \frac{n-1}{r} (1+Br).$$

Then

$$(14) \quad \Delta_2 w \leq \frac{4}{q+1-p} (R^2 - r^2)^{-\frac{2(q+2-p)}{q+1-p}} \left( \frac{2r^2(q+3-p)}{q+1-p} + (R^2 - r^2)(1+(n-1)(1+Br)) \right).$$

Moverover from [3, Chap 2, p. 23]

$$(15) \quad D^2 w = w'' dr \otimes dr + w' D^2 r.$$

If  $0 \geq \text{Sec}_g(x) \geq -S^2$ , there holds

$$(16) \quad 0 \leq D^2 r \leq S \coth(Sr) g \leq \frac{S}{r} (1+Sr) g.$$

Therefore, if  $p \geq 2$  and  $\text{Sec}_g \geq -S^2$ , we get

$$(17) \quad \frac{\langle D^2 w(\nabla u), \nabla u \rangle}{|\nabla u|^2} \leq \frac{4}{q+1-p} (R^2 - r^2)^{-\frac{2(q+2-p)}{q+1-p}} \left( \frac{2r^2(q+3-p)}{q+1-p} + (R^2 - r^2)(2+Sr) \right),$$

while, if  $p \leq 2$  and  $\text{Sec}_g \leq 0$ ,

$$(18) \quad \frac{\langle D^2 w(\nabla u), \nabla u \rangle}{|\nabla u|^2} \leq \frac{4}{q+1-p} (R^2 - r^2)^{-\frac{2(q+2-p)}{q+1-p}} \left( \frac{2r^2(q+3-p)}{q+1-p} + 2(R^2 - r^2) \right).$$

As a consequence

$$(19) \quad \begin{aligned} \mathcal{A}(w) &= -\Delta w - (p-2) \frac{\langle D^2 w(\nabla u), \nabla u \rangle}{|\nabla u|^2} \\ &\geq -k\lambda(R^2 - r^2)^{-\frac{2(q+2-p)}{q+1-p}} (R^2 + (R^2 - r^2)B_p r) \end{aligned}$$

for some  $k = k(n, p, q)$ , where  $B_p = B + (p-2)_+ S$ . Since

$$w^{q+2-p} \geq \lambda^{q+2-p} (R^2 - r^2)^{-\frac{2(q+1-p)}{q+1-p}} + \mu^{q+2-p},$$

we have

$$(20) \quad \begin{aligned} \mathcal{L}^*(w) &\geq \lambda(R^2 - r^2)^{-\frac{2(q+2-p)}{q+1-p}} \left( -k(R^2 + (R^2 - r^2)B_p r) - D \frac{16}{(q+1-p)^2} r^2 + C\lambda^{q+1-p} \right) \\ &\quad + \mu^{q+2-p} - (n-1)B^2\lambda(R^2 - r^2)^{-\frac{2}{q+1-p}} - (n-1)B^2\mu. \end{aligned}$$

We first take

$$(21) \quad \mu = ((n-1)B^2)^{\frac{1}{q+1-p}}.$$

Next we choose  $\lambda$  in order to have, uniformly for  $0 \leq r < R$ ,

$$2^{-1}C\lambda^{q+1-p} \geq k(R^2 + (R^2 - r^2)B_p r) + \frac{16Dr^2}{(q+1-p)^2}$$

and

$$2^{-1}C\lambda^{q+2-p}(R^2 - r^2)^{-\frac{2(q+2-p)}{q+1-p}} \geq (n-1)B^2\lambda(R^2 - r^2)^{-\frac{2}{q+1-p}}.$$

There exists  $c = c(n, p, q)$  such that, if

$$(22) \quad \lambda = c \max \left\{ (R^4 B^2)^{\frac{1}{q+1-p}}, ((1+B_p)R^3)^{\frac{1}{q+1-p}} \right\},$$

then  $\mathcal{L}^*(w) \geq 0$  holds.

**Lemma 2.** *Under the assumptions of Lemma 1, any  $C^1$  solution of (2) in  $M$  satisfies*

$$(23) \quad |\nabla u(x)| \leq c_{n,p,q} \max \left\{ B^{\frac{1}{q+1-p}}, (1+B_p)^{\frac{1}{2(q+1-p)}} (d(x, \partial\Omega))^{-\frac{1}{2(q+1-p)}} \right\} \quad \forall x \in \Omega,$$

for every domain  $\Omega \subset M$ , where  $B_p = B + (p-2)_+ S$  and  $S = S_{d(x, \partial\Omega)}$ .

**Proof.** Assume  $a \in \Omega$ , with  $R < d(a, \partial\Omega)$ . Let  $w$  be as in Lemma 1, then in any connected component  $G$  of  $\{x \in B_R(a) : z(x) - w(x) > 0\}$  we find

$$(24) \quad \mathcal{A}(z-w) + C(z^{q+2-p} - w^{q+2-p}) - (n-1)B^2(z-w) - D \left( \frac{|\nabla z|^2}{z} - \frac{|\nabla w|^2}{w} \right) \leq 0.$$

By the mean value theorem and since  $w(a)$  is the minimum of  $w$ , there holds

$$(25) \quad C(z^{q+2-p} - w^{q+2-p}) - (n-1)B^2(z-w) > 0,$$

provided  $C(q+2-p)(w(a))^{q+1-p} > (n-1)B^2$ . Since  $w(a) > \mu = ((n-1)B^2)^{\frac{1}{q+1-p}}$  and  $q+2-p > 1$ , this condition is fulfilled, up to replacing  $\mu$  by  $A\mu$  for some  $A = A(p, q) > 1$ . If  $x_0 \in G$  is such that  $z - w$  is maximal at  $x_0$ , we derive that

$$\mathcal{A}(z - w) + C(z^{q+2-p} - w^{q+2-p}) - (N-1)B^2(z - w) - D\left(\frac{|\nabla z|^2}{z} - \frac{|\nabla w|^2}{w}\right) \leq 0$$

if  $x = x_0$ , which is a contradiction. Thus  $G = \emptyset$ ,  $z \leq w$  and (23) follows.

The proof of Theorem 1 and Corollary 2 follows by taking  $\Omega = B_R(x)$  and letting  $R \rightarrow \infty$ .

**Proof of Theorem 3.** We take  $q = p$  and assume that  $v$  is  $p$ -harmonic and positive. If we write  $v = e^{-\frac{u}{p-1}}$ , then  $u$  satisfies

$$-\Delta_p u + |\nabla u|^p = 0.$$

If  $\text{Ric}_g(x) \geq 0$ ,  $u$  is constant by Corollary 2, and so is  $v$ . If  $\inf\{\text{Ric}_g(x) : x \in M\} = (1-n)B^2 < 0$  we apply (23) to  $\nabla u$ . If  $\gamma$  is a minimizing geodesic from  $a$  to  $x$ , then  $|\gamma'(t)| = 1$  and

$$u(x) - u(a) = \int_0^{d(x,a)} \frac{d}{dt} u \circ \gamma(t) dt = \int_0^{d(x,a)} \langle \nabla u \circ \gamma(t), \gamma'(t) \rangle dt.$$

Since

$$|\langle \nabla u \circ \gamma(t), \gamma'(t) \rangle| \leq |\nabla u \circ \gamma(t)| \leq c_{n,p,p} B,$$

we obtain

$$(26) \quad u(a) - c_{n,p,p} B \text{dist}(x, a) \leq u(x) \leq u(a) + c_{n,p,p} B \text{dist}(x, a) \quad \forall x \in M.$$

Then (6) follows since  $u = (1-p) \ln v$ .

#### RÉFÉRENCES

- [1] S. Y. Cheng and S.-T. Yau : *Differential equations on Riemannian manifolds and their geometric applications*, **Comm. Pure Appl. Math.** **28**, 333–354 (1975).
- [2] T. H. Colding and W.P. Minicozzi : *An excursion into geometric analysis*, Grigor'yan, Alexander (ed.) et al., *Surveys in differential geometry*. Somerville, MA : International Press, **Surveys in Differential Geometry** **9**, 83–146 (2004).
- [3] R.E. Greene and H. Wu/ *Function Theory on Manifolds Which Possess a Pole*, **Lecture Notes in Mathematics** **699**, Springer-Verlag Berlin Heidelberg (1979).
- [4] B. L. Kotschwar and L. Ni : *Local gradient estimates of  $p$ -harmonic functions,  $1/H$ -flow, and an entropy formula*, **Ann. Sci. École. Norm. Sup.** **42**, 1–36 (2009).
- [5] A. Porretta and L. Véron : *Separable  $p$ -harmonic functions in a cone and related quasilinear equations on manifolds*, **J. Eur. Math. Soc.** **11**, 1285–1305 (2009).
- [6] A. Ratto, M. Rigoli and L. Véron *Conformal immersion of complete Riemannian manifolds and extensions of the Schwarz lemma*, **Duke Math. J.** **74**, 223–236 (1994).